

Double- β Decay

OMMITTED FROM SUMMARY TABLE
A REVIEW GOES HERE – Check our WWW List of Reviews

Half-life Measurements and Limits for Double- β Decay

In most cases the transitions $(Z, A) \rightarrow (Z+2, A) + 2e^- + (0 \text{ or } 2) \bar{\nu}_e$ to the 0^+ ground state of the final nucleus are listed. However, we also list transitions that increase the nuclear charge ($2e^+, e^+/EC$ and $ECEC$) and transitions to excited states of the final nuclei ($0_i^+, 2^+, \text{ and } 2_i^+$). In the following Listings, only best or comparable limits or lifetimes for each isotope are reported and only those with $T_{1/2} > 10^{20}$ years that are relevant for particle physics. For 2ν decay, which is well established, only measured half-lives are reported.

$t_{1/2}(10^{21} \text{ yr})$	CL%	ISOTOPE	TRANSITION	METHOD	DOCUMENT ID	NODE
• • • We do not use the following data for averages, fits, limits, etc. • • •						
>940	90	^{130}Te	0ν	$0^+ \rightarrow 0_1^+$	CUORICINO	1 ANDREOTTI 12
>16000	90	^{136}Xe	0ν	$g.s. \rightarrow g.s.$	EXO-200	2 AUGER 12
> 1.0	90	^{106}Cd	0ν	ECEC, g.s.	$^{106}\text{CdWO}_4$ scint.	3 BELLI 12A
> 2.2	90	^{106}Cd	0ν	EC, β^+ ,g.s.	$^{106}\text{CdWO}_4$ scint.	4 BELLI 12A
> 1.2	90	^{106}Cd	0ν	$\beta^+\beta^+$,g.s.	$^{106}\text{CdWO}_4$ scint.	5 BELLI 12A
$2.38 \pm 0.02 \pm 0.14$	68	^{136}Xe	2ν	$g.s. \rightarrow g.s.$	KamLAND-Zen	6 GANDO 12A
>5700	90	^{136}Xe	0ν	$g.s. \rightarrow g.s.$	KamLAND-Zen	7 GANDO 12A
$2.11 \pm 0.04 \pm 0.21$	68	^{136}Xe	2ν		EXO-200	8 ACKERMAN 11
$0.7 \pm 0.09 \pm 0.11$	68	^{130}Te	2ν		NEMO-3	9 ARNOLD 11
> 130	90	^{130}Te	0ν		NEMO-3	10 ARNOLD 11
> 1.3	90	^{112}Sn	0ν	$0^+ \rightarrow 0_3^+$	γ Ge det.	11 BARABASH 11
> 0.69	90	^{112}Sn	0ν	$0^+ \rightarrow 0_2^+$	γ Ge det.	12 BARABASH 11
> 1.3	90	^{112}Sn	0ν	$0^+ \rightarrow 0_1^+$	γ Ge det.	13 BARABASH 11
> 1.06	90	^{112}Sn	0ν		γ Ge det.	14 BARABASH 11
($2.8 \pm 0.1 \pm 0.3$)E-2		^{116}Cd	2ν		NEMO-3	15 BARABASH 11A
($4.4^{+0.5}_{-0.4} \pm 0.4$)E-2		^{48}Ca	2ν		NEMO-3	16,17 BARABASH 11A
($69 \pm 9 \pm 10$)E-2		^{130}Te	2ν		NEMO-3	17,18 BARABASH 11A
>1100	90	^{100}Mo	0ν		NEMO-3	17,19 BARABASH 11A
>360	90	^{82}Se	0ν		NEMO-3	17,20 BARABASH 11A
>100	90	^{130}Te	0ν		NEMO-3	17,21 BARABASH 11A
>16	90	^{116}Cd	0ν		NEMO-3	17,22 BARABASH 11A
>13	90	^{48}Ca	0ν		NEMO-3	17,23 BARABASH 11A
> 0.32	90	^{64}Zn	0ν	ECEC,g.s.	ZnWO_4 scint.	24 BELLI 11D
> 0.85	90	^{64}Zn	0ν	$\beta^+EC,g.s.$	ZnWO_4 scint.	24 BELLI 11D
> 0.11	90	^{106}Cd	0ν	$0^+ \rightarrow 4^+$	TGV2 det.	25 RUKHADZE 11
($2.35 \pm 0.14 \pm 0.16$)E-2		^{96}Zr	2ν		NEMO-3	26 ARGYRIADES 10
> 9.2	90	^{96}Zr	0ν		NEMO-3	27 ARGYRIADES 10
> 0.22	90	^{96}Zr	0ν	$0^+ \rightarrow 0_1^+$	NEMO-3	28 ARGYRIADES 10
$0.69^{+0.10}_{-0.08} \pm 0.07$	68	^{100}Mo	2ν	$0^+ \rightarrow 0_1^+$	Ge coinc.	29 BELLI 10
> 18.0	90	^{150}Nd	0ν		NEMO-3	30 ARGYRIADES 09
($9.11^{+0.25}_{-0.22} \pm 0.63$)E-3		^{150}Nd	2ν		NEMO-3	31 ARGYRIADES 09
> 0.43	90	^{64}Zn	0ν	β^+EC	ZnWO_4 scint.	32 BELLI 09A
> 0.11	90	^{64}Zn	0ν	ECEC	ZnWO_4 scint.	33 BELLI 09A
$0.55^{+0.12}_{-0.09}$		^{100}Mo	$2\nu+0\nu$	$0^+ \rightarrow 0_1^+$	Ge coincidence	34 KIDD 09
> 3000	90	^{130}Te	0ν		TeO_2 bolometer	35 ARNABOLDI 08
> 0.22	90	^{64}Zn	0ν		ZnWO_4 scint.	36 BELLI 08
> 1.1	90	^{114}Cd	0ν	$\beta\beta$	CdWO_4 scint.	37 BELLI 08B
> 58	90	^{48}Ca	0ν		CaF_2 scint.	38 UMEHARA 08
$0.57^{+0.13}_{-0.09} \pm 0.08$	68	^{100}Mo	2ν	$0^+ \rightarrow 0_1^+$	NEMO-3	39 ARNOLD 07
> 89	90	^{100}Mo	0ν	$0^+ \rightarrow 0_1^+$	NEMO-3	40 ARNOLD 07
> 160	90	^{100}Mo	0ν	$0^+ \rightarrow 2^+$	NEMO-3	41 ARNOLD 07
> 0.0019	90	^{74}Se	$0\nu+2\nu$		γ in Ge det.	42 BARABASH 07
> 0.0055	90	^{74}Se	$0\nu+2\nu$	$0^+ \rightarrow 2_2^+$	γ in Ge det.	43 BARABASH 07
22300^{+4400}_{-3100}	68	^{76}Ge	0ν		Enriched HPGe	44 KLAPDOR-K... 06A
> 1800	90	^{130}Te	0ν		Cryog. det.	45 ARNABOLDI 05
> 460	90	^{100}Mo	0ν		NEMO-3	46 ARNOLD 05A

> 100	90	82Se	0ν	NEMO-3	47	ARNOLD	05A	OCCUR=2	
(7.11 ± 0.02 ± 0.54)E-3	100Mo	2ν		NEMO-3	48	ARNOLD	05A	OCCUR=3	
(9.6 ± 0.3 ± 1.0)E-2	82Se	2ν		NEMO-3	49	ARNOLD	05A	OCCUR=4	
> 140	90	82Se	0ν	NEMO-3	50	ARNOLD	04	OCCUR=2	
(7.68 ± 0.02 ± 0.54)E-3	100Mo	2ν		NEMO-3	51	ARNOLD	04	OCCUR=3	
0.14 ^{+0.04} _{-0.02} ± 0.03	68	150Nd	0ν+2ν	0 ⁺ → 0 ₁ ⁺	γ in Ge det.	52	BARABASH	04	
> 31	90	130Te	0ν	0 ⁺ → 2 ⁺	Cryog. det.	53	ARNABOLDI	03	OCCUR=2
0.61 ± 0.14 ^{+0.29} _{-0.35}	90	130Te	2ν		Cryog. det.	54	ARNABOLDI	03	OCCUR=3
> 110	90	128Te	0ν		Cryog. det.	55	ARNABOLDI	03	OCCUR=4
(0.029 ^{+0.004} _{-0.003})	116Cd	2ν		116CdWO ₄ scint.	56	DANEVICH	03		
> 170	90	116Cd	0ν	116CdWO ₄ scint.	57	DANEVICH	03	OCCUR=2	
> 29	90	116Cd	0ν	0 ⁺ → 2 ⁺	116CdWO ₄ scint.	58	DANEVICH	03	OCCUR=3
> 14	90	116Cd	0ν	0 ⁺ → 0 ₁ ⁺	116CdWO ₄ scint.	59	DANEVICH	03	OCCUR=4
> 6	90	116Cd	0ν	0 ⁺ → 0 ₂ ⁺	116CdWO ₄ scint.	60	DANEVICH	03	OCCUR=5
> 1.1	90	186W	0ν		CdWO ₄ scint.	61	DANEVICH	03	OCCUR=6
> 1.1	90	186W	0ν	0 ⁺ → 2 ⁺	CdWO ₄ scint.	62	DANEVICH	03	OCCUR=7
1.74 ± 0.01 ^{+0.18} _{-0.16}	76Ge	2ν		Enriched HPGe	63	DOERR	03		
>15700	90	76Ge	0ν		Enriched HPGe	64	AALSETH	02B	
> 58	90	134Xe	0ν		Liquid Xe Scint.	65	BERNABEI	02D	
> 1200	90	136Xe	0ν		Liquid Xe Scint.	66	BERNABEI	02D	OCCUR=2
> 4.9	90	100Mo	0ν		Liq. Ar ioniz.	67	ASHITKOV	01	OCCUR=2
> 1.3	90	160Gd	0ν		Gd ₂ SiO ₅ :Ce	68	DANEVICH	01	
> 1.3	90	160Gd	0ν	0 ⁺ → 2 ⁺	Gd ₂ SiO ₅ :Ce	69	DANEVICH	01	OCCUR=2
0.59 ^{+0.17} _{-0.11} ± 0.06	100Mo	0ν+2ν	0 ⁺ → 0 ₁ ⁺	Ge coinc.	70	DEBRAECKEL.01			
>19000	90	76Ge	0ν		Enriched HPGe	71	KLAPDOR-K...01		
(9.4 ± 3.2)E-3	90	96Zr	0ν+2ν		Geochem	72	WIESER	01	
0.042 ^{+0.033} _{-0.013}	48Ca	2ν		Ge spectrometer	73	BRUDANIN	00	OCCUR=2	
0.021 ^{+0.008} _{-0.004} ± 0.002	96Zr	2ν		NEMO-2	74	ARNOLD	99		
(8.3 ± 1.0 ± 0.7)E-2	82Se	2ν		NEMO-2	75	ARNOLD	98		
> 2.8	90	82Se	0ν	0 ⁺ → 2 ⁺	NEMO-2	76	ARNOLD	98	OCCUR=3
(7.6 ^{+2.2} _{-1.4})E-3	100Mo	2ν		Si(Li)	77	ALSTON-...	97	OCCUR=2	
(6.82 ^{+0.38} _{-0.53} ± 0.68)E-3	100Mo	2ν		TPC	78	DESilva	97		
(6.75 ^{+0.37} _{-0.42} ± 0.68)E-3	150Nd	2ν		TPC	79	DESilva	97	OCCUR=3	
(3.75 ± 0.35 ± 0.21)E-2	116Cd	2ν	0 ⁺ → 0 ⁺	NEMO 2	80	ARNOLD	96		
0.043 ^{+0.024} _{-0.011} ± 0.014	48Ca	2ν		TPC	81	BALYSH	96		
0.79 ± 0.10	130Te	0ν+2ν		Geochem	82	TAKAOKA	96		
0.61 ^{+0.18} _{-0.11}	100Mo	0ν+2ν	0 ⁺ → 0 ₁ ⁺	Geochem	83	BARABASH	95		
0.026 ^{+0.009} _{-0.005}	116Cd	2ν	0 ⁺ → 0 ⁺	ELEGANT IV	EJIRI	95			
0.017 ^{+0.010} _{-0.005} ± 0.0035	150Nd	2ν	0 ⁺ → 0 ⁺	TPC	ARTEMEV	93			
0.039 ± 0.009	96Zr	0ν+2ν		Geochem	KAWASHIMA	93			
2.7 ± 0.1	130Te	0ν+2ν		Geochem	BERNATOW...	92			
7200 ± 400	128Te	0ν+2ν		Geochem	84	BERNATOW...	92	OCCUR=2	
0.108 ^{+0.026} _{-0.006}	82Se	2ν	0 ⁺ → 0 ⁺	TPC	ELLIOTT	92		OCCUR=2	
2.0 ± 0.6	238U	0ν+2ν		Radiochem	85	TURKEVICH	91		
0.12 ± 0.01 ± 0.04	68	82Se	0ν+2ν		86 LIN	88			
0.75 ± 0.03 ± 0.23	68	130Te	0ν+2ν		87 LIN	88	OCCUR=2		
1800 ± 700	68	128Te	0ν+2ν		88 LIN	88B			
2.60 ± 0.28	130Te	0ν+2ν		Geochem	89 KIRSTEN	83		OCCUR=2	

1 ANDREOTTI 12 use high resolution TeO₂ bolometric calorimeter to search for the 0νββ decay of ¹³⁰Te leading to the excited 0₊¹ state at 1793.5 keV.

2 AUGER 12 use EXO-200 liquid Xe TPC filled with 79.4 kg (fiducial mass) of ¹³⁶Xe to constrain the 0νββ-decay halflife of ¹³⁶Xe. This result is more sensitive than BERN-ABEI 02D and GANDO 12A.

3 BELLINI 12A use ¹⁰⁶CdWO₄ 215 g crystal scintillator to search for various ββ decay modes. The limit for the ECEC mode is derived from the fit to the background spectrum in the 1.8–3.2 MeV energy interval in the run of 6590 hours. The same analysis provides several limits (~ 2–5 × 10²⁰ years) for the EEC mode leading to the excited 0⁺ and 2⁺ states. Also a similar size limits for the possible resonance process populating states at 2718 keV, 2741 keV, and 2748 keV were obtained.

NODE=S076HDB;LINKAGE=ND

NODE=S076HDB;LINKAGE=AU

NODE=S076HDB;LINKAGE=IB

- ⁴ BELLI 12A use $^{106}\text{CdWO}_4$ 215 g crystal scintillator to search for various $\beta\beta$ decay modes. The limit for the $\text{EC}\beta^+$ mode is derived from the fit to the background spectrum in the 2.0–3.0 MeV energy interval in the run of 6590 hours. The same analysis provides several limits ($\sim 0.5\text{--}1.3 \times 10^{21}$ years) for the $\text{EC}\beta^+$ mode leading to the excited 0^+ and 2^+ states.
- ⁵ BELLI 12A use $^{106}\text{CdWO}_4$ 215 g crystal scintillator to search for various $\beta\beta$ decay modes. The limit for the $\beta^+\beta^+$ mode is derived from the fit to the background spectrum in the 0.76–2.8 MeV energy interval in the run of 6590 hours. The same analysis provides the limit (1.2×10^{21} years) for the $\beta^+\beta^+$ mode leading to the first excited 2^+ state.
- ⁶ GANDO 12A use a modification of the existing KamLAND detector. The $\beta\beta$ decay source/detector is 13 tons of enriched ^{136}Xe -loaded scintillator contained in an inner balloon. The $2\nu\beta\beta$ decay rate is derived from the fit to the spectrum between 0.5 and 4.8 MeV. This result is in agreement with ACKERMAN 11.
- ⁷ GANDO 12A use a modification of the existing KamLAND detector. The $\beta\beta$ decay source/detector is 13 tons of enriched ^{136}Xe -loaded scintillator contained in an inner balloon. The $0\nu\beta\beta$ decay rate is derived from the fit where the background rates were allowed to float. This result is more sensitive than BERNABEI 02D.
- ⁸ ACKERMAN 11 use the EXO-200 liquid Xe TPC filled with ~ 175 kg of enriched ^{136}Xe to determine the 2ν halflife of ^{136}Xe .
- ⁹ ARNOLD 11 use enriched ^{130}Te in the NEMO-3 detector to measure the $2\nu\beta\beta$ decay rate. This result is in agreement with, but more accurate than ARNABOLDI 03.
- ¹⁰ ARNOLD 11 use the NEMO-3 detector to obtain a limit for the $0\nu\beta\beta$ decay. This result is less significant than ARNABOLDI 05.
- ¹¹ BARABASH 11 use 100 g of enriched ^{112}Sn to determine a limit for the ECEC 0ν decay to the 0_3^+ state of ^{112}Cd by searching for the de-excitation γ with a Ge detector. This decay mode is a candidate for resonant rate enhancement.
- ¹² BARABASH 11 use 100 g of enriched ^{112}Sn to determine a limit for the ECEC 0ν decay to the 0_2^+ state of ^{112}Cd by searching for the de-excitation γ with a Ge detector.
- ¹³ BARABASH 11 use 100 g of enriched ^{112}Sn to determine a limit for the ECEC 0ν decay to the 0_1^+ state of ^{112}Cd by searching for the de-excitation γ with a Ge detector.
- ¹⁴ BARABASH 11 use 100 g of enriched ^{112}Sn to determine a limit for the ECEC 0ν decay to the ground state of ^{112}Cd by searching for the de-excitation γ with a Ge detector.
- ¹⁵ Supersedes DANEVICH 03 and ARNOLD 96.
- ¹⁶ Supersedes BRUDANIN 00 and BALYSH 96.
- ¹⁷ BARABASH 11A use the NEMO-3 detector to measure $\beta\beta 2\nu$ rates and place limits on $\beta\beta 0\nu$ half lives for various nuclides.
- ¹⁸ Supersedes ARNABOLDI 03.
- ¹⁹ Supersedes ARNOLD 05A, ARNOLD 04, ASHITKOV 01, EJIRI 01, and DASSIE 95.
- ²⁰ Supersedes ARNOLD 05A, ARNOLD 04, ARNOLD 98, and ELLIOTT 92.
- ²¹ Less restrictive than ARNABOLDI 08.
- ²² Less restrictive than DANEVICH 03.
- ²³ Less restrictive than UMEHARA 08 and OGAWA 04.
- ²⁴ BELLI 11D use ZnWO_4 scintillator calorimeters to search for various $\beta\beta$ decay modes of ^{64}Zn , ^{70}Zn , ^{180}W , and ^{186}W .
- ²⁵ RUKHADZE 11 uses 13.6 g of enriched ^{106}Cd to search for the neutrinoless ECEC decay into an excited state of ^{106}Pd and its characteristic γ -radiation using the TGV2 detector. This decay mode is a candidate for resonant rate enhancement, however, hindered by the large spin difference.
- ²⁶ ARGYRIADES 10 use 9.4 ± 0.2 g of ^{96}Zr in NEMO-3 detector and identify its $2\nu\beta\beta$ decay. The result is in agreement and supersedes ARNOLD 99.
- ²⁷ ARGYRIADES 10 use 9.4 ± 0.2 g of ^{96}Zr in NEMO-3 detector and obtain a limit of the $0\nu\beta\beta$ decay. The result is in agreement and supersedes ARNOLD 99.
- ²⁸ ARGYRIADES 10 use 9.4 ± 0.2 g of ^{96}Zr in NEMO-3 detector and obtain a limit of the $0\nu\beta\beta$ decay into the first excited 0_1^+ state in ^{96}Mo .
- ²⁹ BELLI 10 use enriched ^{100}Mo with 4 HP Ge detectors to record the 590.8 and 539.5 keV γ rays from the decay of the 0_1^+ state in ^{100}Ru both in singles and coincidences. This result confirms the measurement of KIDD 09 and ARNOLD 07 and supersedes them.
- ³⁰ ARGYRIADES 09 use the NEMO-3 tracking calorimeter containing 36.5 g of ^{150}Nd , a total exposure of 924.7 days, to derive a limit for the $0\nu\beta\beta$ half-life. Supersedes DESILVA 97.
- ³¹ ARGYRIADES 09 use the NEMO-3 tracking calorimeter containing 36.5 g of ^{150}Nd , a total exposure of 924.7 days, to determine the value of the $2\nu\beta\beta$ half-life. This result is in marginal agreement, but has somewhat smaller error bars, than DESILVA 97.
- ³² BELLI 09A use ZnWO_4 scintillating crystals to search for various modes of $\beta\beta$ decay. This work improves the limits for different modes of ^{64}Zn decay into the ground state of ^{64}Ni , in this case for the $0\nu\beta^+\text{EC}$ mode. Supersedes BELLI 08.
- ³³ BELLI 09A use ZnWO_4 scintillating crystals to search for various modes of $\beta\beta$ decay. This work improves the limits for different modes of ^{64}Zn decay into the ground state of ^{64}Ni , in this case for the $0\nu\text{ECEC}$ mode. Supersedes BELLI 08.

NODE=S076HDB;LINKAGE=IL

NODE=S076HDB;LINKAGE=PB

NODE=S076HDB;LINKAGE=GA

NODE=S076HDB;LINKAGE=GN

NODE=S076HDB;LINKAGE=CK

NODE=S076HDB;LINKAGE=AD

NODE=S076HDB;LINKAGE=RD

NODE=S076HDB;LINKAGE=B3

NODE=S076HDB;LINKAGE=B2

NODE=S076HDB;LINKAGE=B4

NODE=S076HDB;LINKAGE=B0

NODE=S076HDB;LINKAGE=SB

NODE=S076HDB;LINKAGE=H2

NODE=S076HDB;LINKAGE=SA

NODE=S076HDB;LINKAGE=H3

NODE=S076HDB;LINKAGE=H4

NODE=S076HDB;LINKAGE=H5

NODE=S076HDB;LINKAGE=H6

NODE=S076HDB;LINKAGE=H7

NODE=S076HDB;LINKAGE=H8

NODE=S076HDB;LINKAGE=LE

NODE=S076HDB;LINKAGE=RU

NODE=S076HDB;LINKAGE=RG

NODE=S076HDB;LINKAGE=RY

NODE=S076HDB;LINKAGE=RR

NODE=S076HDB;LINKAGE=IE

NODE=S076HDB;LINKAGE=AG

NODE=S076HDB;LINKAGE=AY

NODE=S076HDB;LINKAGE=LB

NODE=S076HDB;LINKAGE=LI

- 34 KIDD 09 combine past and new data with an improved coincidence detection efficiency determination. The result agrees with ARNOLD 95. Supersedes DEBRAECKELEER 01 and BARABASH 95.
- 35 ARNABOLDI 08 use high resolution TeO_2 bolometer calorimeter to search for double beta decay of ^{130}Te . Supersedes ARNABOLDI 05.
- 36 BELLI 08 use ZnWO_4 scintillation calorimeter to search for neutrinoless β^+ plus electron capture decay of ^{64}Zn . The halflife limit for the 2ν mode is 2.1×10^{20} years.
- 37 BELLI 08B use CdWO_4 scintillation calorimeter to search for $0\nu\beta\beta$ decay of ^{114}Cd .
- 38 UMEHARA 08 use CaF_2 scintillation calorimeter to search for double beta decay of ^{48}Ca . Limit is significantly more stringent than quoted sensitivity: 18×10^{21} years.
- 39 First exclusive measurement of 2ν -decay to the first excited 0_1^+ -state of daughter nucleus. ARNOLD 07 use the NEMO-3 tracking calorimeter to detect all particles emitted in decay. Result agrees with the inclusive ($0\nu + 2\nu$) measurement of DEBRAECKELEER 01.
- 40 Limit on 0ν -decay to the first excited 0_1^+ -state of daughter nucleus using NEMO-3 tracking calorimeter. Supersedes DASSIE 95.
- 41 Limit on 0ν -decay to the first excited 2^+ -state of daughter nucleus using NEMO-3 tracking calorimeter.
- 42 BARABASH 07 use Ge calorimeter to search for γ -radiation following double electron capture or β^+ plus electron capture decays of ^{74}Se to the ground state of ^{74}Ge . This limit is based on the search for the 511 keV annihilation radiation. Various other limits, for the capture from different atomic shells and also to the excited states, are reported in the paper.
- 43 BARABASH 07 use Ge calorimeter to search for γ -radiation following double electron capture decay of ^{74}Se into the second excited 2^+ -state of ^{74}Ge . That transition has been considered due to a possible resonance enhancement. The 2ν mode would be suppressed for this decay by its extremely small phase space factor.
- 44 Klapdor-Kleingrothaus 06A present re-analysis of data originally published in Klapdor-Kleingrothaus 04A. Modified pulse shape analysis leads the authors to claim improved 6σ statistical evidence for observation of 0ν -decay, compared to 4.2σ in Klapdor-Kleingrothaus 04A. Analysis of the systematic uncertainty is not presented. Supersedes Klapdor-Kleingrothaus 04A.
- 45 Supersedes ARNABOLDI 04. Bolometric TeO_2 detector array CUORICINO is used for high resolution search for $0\nu\beta\beta$ decay. The half-life limit is derived from 3.09 kg yr ^{130}Te exposure.
- 46 NEMO-3 tracking calorimeter containing 6.9 kg of enriched ^{100}Mo is used in ARNOLD 05A. A limit for $0\nu\beta\beta$ half-life of ^{100}Mo is reported. Supersedes ARNOLD 04.
- 47 NEMO-3 tracking calorimeter is used in ARNOLD 05A to place limit on $0\nu\beta\beta$ half-life of ^{82}Se . Detector contains 0.93 kg of enriched ^{82}Se . Supersedes ARNOLD 04.
- 48 ARNOLD 05A use the NEMO-3 tracking calorimeter to determine the $2\nu\beta\beta$ half-life of ^{100}Mo with high statistics and low background (389 days of data taking). Supersedes ARNOLD 04.
- 49 ARNOLD 05A use the NEMO-3 tracking detector to determine the $2\nu\beta\beta$ half-life of ^{82}Se with high statistics and low background (389 days of data taking). Supersedes ARNOLD 04.
- 50 ARNOLD 04 use the NEMO-3 tracking detector to determine the limit for $0\nu\beta\beta$ halflife of ^{82}Se . This represents an improvement, by a factor of ~ 10 , when compared with ELLIOTT 92. It supersedes the limit of ARNOLD 98 for this decay using NEMO-2.
- 51 ARNOLD 04 use the NEMO-3 tracking detector to determine the $2\nu\beta\beta$ halflife of ^{100}Mo with high statistics and low background. The halflife is determined assuming the Single State Dominance. It is in agreement with, and more accurate than, previous determinations. Supersedes DASSIE 95 determination of this quantity with NEMO-2.
- 52 BARABASH 04 perform an inclusive measurement of the $\beta\beta$ decay of ^{150}Nd into the first excited (0_1^+) state of ^{150}Sm . Gamma radiation emitted in decay of the excited state is detected.
- 53 Decay into first excited state of daughter nucleus.
- 54 Two neutrino decay into ground state. Relatively large error mainly due to uncertainties in background determination. Reported value is shorter than the geochemical measurements of KIRSTEN 83 and BERNATOWICZ 92 but in agreement with LIN 88 and TAKAOKA 96.
- 55 Supersedes ALESSANDRELLO 00. Array of TeO_2 crystals in high resolution cryogenic calorimeter. Some enriched in ^{128}Te . Ground state to ground state decay.
- 56 Calorimetric measurement of 2ν ground state decay of ^{116}Cd using enriched CdWO_4 scintillators. Agrees with EJIRI 95 and ARNOLD 96. Supersedes DANEVICH 00.
- 57 Limit on 0ν decay of ^{116}Cd using enriched CdWO_4 scintillators. Supersedes DANEVICH 00.
- 58 Limit on 0ν decay of ^{116}Cd into first excited 2^+ state of daughter nucleus using enriched CdWO_4 scintillators. Supersedes DANEVICH 00.
- 59 Limit on 0ν decay of ^{116}Cd into first excited 0^+ state of daughter nucleus using enriched CdWO_4 scintillators. Supersedes DANEVICH 00.
- 60 Limit on 0ν decay of ^{116}Cd into second excited 0^+ state of daughter nucleus using enriched CdWO_4 scintillators. Supersedes DANEVICH 00.
- 61 Limit on the 0ν ground state decay of ^{186}W using enriched CdWO_4 scintillators.

- 62 Limit on the 0ν decay of ^{186}W to the first excited 2^+ state of the daughter nucleus using enriched CdWO_4 scintillators.
- 63 Results of the Heidelberg-Moscow experiment (KLAUDOR-KLEINGROTHAUS 01 and GUENTHER 97) are reanalyzed using a new simulation of the complete background spectrum. The $\beta\beta 2\nu$ -decay rate is deduced from a 41.57 kg-y exposure. The result is in agreement and supersedes the above referenced halflives with similar statistical and systematic errors.
- 64 AALSETH 02B limit is based on 117 mol-yr of data using enriched Ge detectors. Background reduction by means of pulse shape analysis is applied to part of the data set. Reported limit is slightly less restrictive than that in KLAUDOR-KLEINGROTHAUS 01. However, it excludes part of the allowed half-life range reported in KLAUDOR-KLEINGROTHAUS 01B for the same nuclide. The analysis has been criticized in KLAUDOR-KLEINGROTHAUS 04B. The criticism was addressed and disputed in AALSETH 04.
- 65 BERNABEI 02D report a limit for the $0\nu, 0^+ \rightarrow 0^+$ decay of ^{134}Xe , present in the source at 17%, by considering the maximum number of events for this mode compatible with the fitted smooth background.
- 66 BERNABEI 02D report a limit for the $0\nu, 0^+ \rightarrow 0^+$ decay of ^{136}Xe , by considering the maximum number of events for this mode compatible with the fitted smooth background. The quoted sensitivity is 450×10^{21} yr. The Feldman and Cousins method is used to obtain the quoted limit.
- 67 ASHITKOV 01 result for 0ν of ^{100}Mo is less stringent than EJIRI 01.
- 68 DANEVICH 01 place limit on 0ν decay of ^{160}Gd using $\text{Gd}_2\text{SiO}_5:\text{Ce}$ crystal scintillators. The limit is more stringent than KOBAYASHI 95.
- 69 DANEVICH 01 place limits on 0ν decay of ^{160}Gd into excited 2^+ state of daughter nucleus using $\text{Gd}_2\text{SiO}_5:\text{Ce}$ crystal scintillators.
- 70 DEBRAECKEER 01 performed an inclusive measurement of the $\beta\beta$ decay into the second excited state of the daughter nucleus. A novel coincidence technique counting the de-excitation photons is employed. The result agrees with BARABASH 95.
- 71 KLAUDOR-KLEINGROTHAUS 01 is a continuation of the work published in BAUDIS 99. Isotopically enriched Ge detectors are used in calorimetric measurement. The most stringent bound is derived from the data set in which pulse-shape analysis has been used to reduce background. Exposure time is 35.5 kg y. Supersedes BAUDIS 99 as most stringent result.
- 72 WIESER 01 reports an inclusive geochemical measurement of ^{96}Zr $\beta\beta$ half life. Their result agrees within 2σ with ARNOLD 99 but only marginally, within 3σ , with KAWASHIMA 93.
- 73 BRUDANIN 00 determine the 2ν halflife of ^{48}Ca . Their value is less accurate than BALYSH 96.
- 74 ARNOLD 99 measure directly the 2ν decay of Zr for the first time, using the NEMO-2 tracking detector and an isotopically enriched source. The lifetime is more accurate than the geochemical result of KAWASHIMA 93.
- 75 ARNOLD 98 measure the 2ν decay of ^{82}Se by comparing the spectra in an enriched and natural selenium source using the NEMO-2 tracking detector. The measured half-life is in agreement, perhaps slightly shorter, than ELLIOTT 92.
- 76 ARNOLD 98 determine the limit for 0ν decay to the excited 2^+ state of ^{82}Se using the NEMO-2 tracking detector.
- 77 ALSTON-GARNJOST 97 report evidence for 2ν decay of ^{100}Mo . This decay has been also observed by EJIRI 91, DASSIE 95, and DESILVA 97.
- 78 DESILVA 97 result for 2ν decay of ^{100}Mo is in agreement with ALSTON-GARNJOST 97 and DASSIE 95. This measurement has the smallest errors.
- 79 DESILVA 97 result for 2ν decay of ^{150}Nd is in marginal agreement with ARTEMEV 93. It has smaller errors.
- 80 ARNOLD 96 measure the 2ν decay of ^{116}Cd . This result is in agreement with EJIRI 95, but has smaller errors. Supersedes ARNOLD 95.
- 81 BALYSH 96 measure the 2ν decay of ^{48}Ca , using a passive source of enriched ^{48}Ca in a TPC.
- 82 TAKAOKA 96 measure the geochemical half-life of ^{130}Te . Their value is in disagreement with the quoted values of BERNATOWICZ 92 and KIRSTEN 83; but agrees with several other unquoted determinations, e.g., MANUEL 91.
- 83 BARABASH 95 cannot distinguish 0ν and 2ν , but it is inferred indirectly that the 0ν mode accounts for less than 0.026% of their event sample. They also note that their result disagrees with the previous experiment by the NEMO group (BLUM 92).
- 84 BERNATOWICZ 92 finds $^{128}\text{Te}/^{130}\text{Te}$ activity ratio from slope of $^{128}\text{Xe}/^{132}\text{Xe}$ vs $^{130}\text{Xe}/^{132}\text{Xe}$ ratios during extraction, and normalizes to lead-dated ages for the ^{130}Te lifetime. The authors state that their results imply that "(a) the double beta decay of ^{128}Te has been firmly established and its half-life has been determined ... without any ambiguity due to trapped Xe interferences... (b) Theoretical calculations ... underestimate the [long half-lives of ^{128}Te ^{130}Te] by 1 or 2 orders of magnitude, pointing to a real suppression in the 2ν decay rate of these isotopes. (c) Despite [this], most $\beta\beta$ -models predict a ratio of 2ν decay widths ... in fair agreement with observation." Further details of the experiment are given in BERNATOWICZ 93. Our listed half-life has been revised downward from the published value by the authors, on the basis of reevaluated cosmic-ray ^{128}Xe production corrections.
- 85 TURKEVICH 91 observes activity in old U sample. The authors compare their results with theoretical calculations. They state "Using the phase-space factors of Boehm and

NODE=S076HDB;LINKAGE=D7

NODE=S076HDB;LINKAGE=DO

NODE=S076HDB;LINKAGE=SH

NODE=S076HDB;LINKAGE=D5

NODE=S076HDB;LINKAGE=D6

NODE=S076HDB;LINKAGE=S2

NODE=S076HDB;LINKAGE=TH

NODE=S076HDB;LINKAGE=TI

NODE=S076HDB;LINKAGE=RB

NODE=S076HDB;LINKAGE=K1

NODE=S076HDB;LINKAGE=WR

NODE=S076HDB;LINKAGE=P2

NODE=S076HDB;LINKAGE=LD

NODE=S076HDB;LINKAGE=R1

NODE=S076HDB;LINKAGE=R3

NODE=S076HDB;LINKAGE=ZC

NODE=S076HDB;LINKAGE=DA

NODE=S076HDB;LINKAGE=DC

NODE=S076HDB;LINKAGE=AR

NODE=S076HDB;LINKAGE=Y2

NODE=S076HDB;LINKAGE=J3

NODE=S076HDB;LINKAGE=Y1

NODE=S076HDB;LINKAGE=BB

NODE=S076HDB;LINKAGE=S

Vogel (BOEHM 87) leads to matrix element values for the ^{238}U transition in the same range as deduced for ^{130}Te and ^{76}Ge . On the other hand, the latest theoretical estimates (STAUDT 90) give an upper limit that is 10 times lower. This large discrepancy implies either a defect in the calculations or the presence of a faster path than the standard two-neutrino mode in this case." See BOEHM 87 and STAUDT 90.

86 Result agrees with direct determination of ELLIOTT 92.

87 Inclusive half life inferred from mass spectroscopic determination of abundance of $\beta\beta$ -decay product ^{130}Te in mineral kitkaite (NiTeSe). Systematic uncertainty reflects variations in U-Xe gas-retention-age derived from different uranite samples. Agrees with geochemical determination of TAKAOKA 96 and direct measurement of ARNABOLDI 03. Inconsistent with results of KIRSTEN 83 and BERNATOWICZ 92.

88 Ratio of inclusive double beta half lives of ^{128}Te and ^{130}Te determined from minerals melonite (NiTe_2) and altaite (PbTe) by means of mass spectroscopic measurement of abundance of $\beta\beta$ -decay products. As gas-retention-age could not be determined the authors use half life of ^{130}Te (LIN 88) to infer the half life of ^{128}Te . No estimate of the systematic uncertainty of this method is given. The directly determined half life ratio agrees with BERNATOWICZ 92. However, the inferred ^{128}Te half life disagrees with KIRSTEN 83 and BERNATOWICZ 92.

89 KIRSTEN 83 reports "2 σ " error. References are given to earlier determinations of the ^{130}Te lifetime.

NODE=S076HDB;LINKAGE=LN
NODE=S076HDB;LINKAGE=LO

NODE=S076HDB;LINKAGE=LP

NODE=S076HDB;LINKAGE=D

NODE=S076MW

NODE=S076MW

NODE=S076MW

$\langle m_\nu \rangle$, The Effective Weighted Sum of Majorana Neutrino Masses Contributing to Neutrinoless Double- β Decay

$\langle m_\nu \rangle = |\sum U_{1j}^2 m_{\nu_j}|$, where the sum goes from 1 to n and where n = number of neutrino generations, and ν_j is a Majorana neutrino. Note that U_{ej}^2 , not $|U_{ej}|^2$, occurs in the sum. The possibility of cancellations has been stressed. In the following Listings, only best or comparable limits or lifetimes for each isotope are reported.

VALUE (eV)	CL%	ISOTOPE	TRANSITION	METHOD	DOCUMENT ID	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 0.14–0.38	90	^{136}Xe	$0\nu, \text{g.s.} \rightarrow \text{g.s.}$	EXO-200	90 AUGER 12	
< 0.3–0.6	90	^{136}Xe	$0\nu, \text{g.s.} \rightarrow \text{g.s.}$	KamLAND-Zen	91 GANDO 12A	OCCUR=2
< 0.45–0.93	90	^{100}Mo	0ν	NEMO-3	92 BARABASH 11A	OCCUR=2
< 0.89–2.43	90	^{82}Se	0ν	NEMO-3	93 BARABASH 11A	
< 7.2–19.5	90	^{96}Zr	0ν	NEMO-3	94 ARGYRIADES 10	
< 4.0–6.8	90	^{150}Nd	0ν	NEMO-3	95 ARGYRIADES 09	
< 0.19–0.68	90	^{130}Te	0ν	TeO ₂ bolometer	96 ARNABOLDI 08	
< 3.5–22	90	^{48}Ca	0ν	CaF ₂ scint.	97 UMEHARA 08	
< 9.3–60	90	^{100}Mo	$0^+ \rightarrow 0^+_1$	NEMO-3	98 ARNOLD 07	
< 6500	90	^{100}Mo	$0^+ \rightarrow 2^+$	NEMO-3	99 ARNOLD 07	OCCUR=2
0.32±0.03	68	^{76}Ge	0ν	Enriched HPGe	100 Klapdor-K... 06A	
< 0.2–1.1	90	^{130}Te		Cryog. det.	101 ARNABOLDI 05	
< 0.7–2.8	90	^{100}Mo	0ν	NEMO-3	102 ARNOLD 05A	
< 1.7–4.9	90	^{82}Se	0ν	NEMO-3	103 ARNOLD 05A	OCCUR=2
< 0.37–1.9	90	^{130}Te		Cryog. det.	104 ARNABOLDI 04	
< 0.8–1.2	90	^{100}Mo	0ν	NEMO-3	105 ARNOLD 04	
< 1.5–3.1	90	^{82}Se	0ν	NEMO-3	105 ARNOLD 04	OCCUR=2
0.1–0.9	99.7	^{76}Ge		Enriched HP Ge	106 Klapdor-K... 04A	
< 7.2–44.7	90	^{48}Ca		CaF ₂ scint.	107 OGAWA 04	
< 1.1–2.6	90	^{130}Te		Cryog. det.	108 ARNABOLDI 03	
< 1.5–1.7	90	^{116}Cd	0ν	$^{116}\text{CdWO}_4$ scint.	109 DANEVICH 03	
< 0.33–1.35	90			Enriched HPGe	110 AALSETH 02B	
< 2.9	90	^{136}Xe	0ν	Liquid Xe Scint.	111 BERNABEI 02D	
0.39±0.17 –0.28	76	^{76}Ge	0ν	Enriched HPGe	112 Klapdor-K... 02D	
< 2.1–4.8	90	^{100}Mo	0ν	ELEGANT V	113 EJIRI 01	
< 0.35	90	^{76}Ge		Enriched HPGe	114 Klapdor-K... 01	
< 23	90	^{96}Zr		NEMO-2	115 ARNOLD 99	
< 1.1–1.5		^{128}Te		Geochem	116 BERNATOW... 92	
< 5	68	^{82}Se		TPC	117 ELLIOTT 92	
< 8.3	76	^{48}Ca	0ν	CaF ₂ scint.	YOU 91	

90 AUGER 12 limit is based on the EXO-200 data. The reported range reflects different nuclear matrix elements.

NODE=S076MW;LINKAGE=AE

91 GANDO 12A limit is based on the KamLAND-Zen data. The reported range reflects different nuclear matrix elements.

NODE=S076MW;LINKAGE=GA

92 BARABASH 11A limit is based on NEMO-3 data for ^{100}Mo . The reported range reflects different nuclear matrix elements. Supersedes ARNOLD 05A and ARNOLD 04.

NODE=S076MW;LINKAGE=HS

93 BARABASH 11A limit is based on NEMO-3 data for ^{82}Se . The reported range reflects different nuclear matrix elements. Supersedes ARNOLD 05A and ARNOLD 04.

NODE=S076MW;LINKAGE=HA

- 94 ARGYRIADES 10 use ^{96}Zr and the NEMO-3 tracking detector to obtain the reported mass limit. The range reflects the fluctuation of the nuclear matrix elements considered.
- 95 ARGYRIADES 09 limit is based on data taken with the NEMO-3 detector and ^{150}Nd . A range of nuclear matrix elements that include the effect of nuclear deformation have been used.
- 96 Limit was obtained using high resolution TeO_2 bolometer calorimeter to search for double beta decay of ^{130}Te . Reported range of limits reflects spread of matrix element calculations used. Supersedes ARNABOLDI 05.
- 97 Limit was obtained using CaF_2 scintillation calorimeter to search for double beta decay of ^{48}Ca . Reported range of limits reflects spread of QRPA and SM matrix element calculations used. Supersedes OGAWA 04.
- 98 ARNOLD 07 use NEMO-3 half life limit for 0ν -decay of ^{100}Mo to the first excited 0_1^+ -state of daughter nucleus to obtain neutrino mass limit. The spread reflects the choice of two different nuclear matrix elements. This limit is not competitive when compared to the decay to the ground state.
- 99 ARNOLD 07 use NEMO-3 half life limit for 0ν -decay of ^{100}Mo to the first excited 2^+ -state of daughter nucleus to obtain neutrino mass limit. This limit is not competitive when compared to the decay to the ground state.
- 100 Re-analysis of data originally published in Klapdor-Kleingrothaus 04A. Modified pulse shape analysis leads the authors to claim 6σ statistical evidence for observation of 0ν -decay. Authors use matrix element of STAUDT 90. Uncertainty of nuclear matrix element is not reflected in stated error. Supersedes Klapdor-Kleingrothaus 04A.
- 101 Supersedes ARNABOLDI 04. Reported range of limits due to use of different nuclear matrix element calculations.
- 102 Mass limits reported in ARNOLD 05A are derived from ^{100}Mo data, obtained by the NEMO-3 collaboration. The range reflects the spread of matrix element calculations considered in this work. Supersedes ARNOLD 04.
- 103 Neutrino mass limits based on ^{82}Se data utilizing the NEMO-3 detector. The range reported in ARNOLD 05A reflects the spread of matrix element calculations considered in this work. Supersedes ARNOLD 04.
- 104 Supersedes ARNABOLDI 03. Reported range of limits due to use of different nuclear matrix element calculations.
- 105 ARNOLD 04 limit is based on the nuclear matrix elements of SIMKOVIC 99, STOICA 01 and CIVITARESE 03.
- 106 Supersedes Klapdor-Kleingrothaus 02D. Event excess at $\beta\beta$ -decay energy is used to derive Majorana neutrino mass using the nuclear matrix elements of STAUDT 90. The mass range shown is based on the authors evaluation of the uncertainties of the STAUDT 90 matrix element calculation. If this uncertainty is neglected, and only statistical errors are considered, the range in $\langle m \rangle$ becomes (0.2–0.6) eV at the 3σ level.
- 107 Calorimetric CaF_2 scintillator. Range of limits reflects authors' estimate of the uncertainty of the nuclear matrix elements. Replaces YOU 91 as the most stringent limit based on ^{48}Ca .
- 108 Supersedes ALESSANDRELLO 00. Cryogenic calorimeter search. Reported a range reflecting uncertainty in nuclear matrix element calculations.
- 109 Limit for $\langle m_\nu \rangle$ is based on the nuclear matrix elements of STAUDT 90 and ARNOLD 96. Supersedes DANEVICH 00.
- 110 AALSETH 02B reported range of limits on $\langle m_\nu \rangle$ reflects the spread of theoretical nuclear matrix elements. Excludes part of allowed mass range reported in Klapdor-Kleingrothaus 01B.
- 111 BERNABEI 02D limit is based on the matrix elements of SIMKOVIC 02. The range of neutrino masses based on a variety of matrix elements is 1.1–2.9 eV.
- 112 Klapdor-Kleingrothaus 02D is a detailed description of the analysis of the data collected by the Heidelberg-Moscow experiment, previously presented in Klapdor-Kleingrothaus 01B. Matrix elements in STAUDT 90 have been used. See the footnote in the preceding table for further details. See also Klapdor-Kleingrothaus 02B.
- 113 The range of the reported $\langle m_\nu \rangle$ values reflects the spread of the nuclear matrix elements. On axis value assuming $\langle \lambda \rangle = \langle \eta \rangle = 0$.
- 114 Klapdor-Kleingrothaus 01 uses the calculation by STAUDT 90. Using several other models in the literature could worsen the limit up to 1.2 eV. This is the most stringent experimental bound on m_ν . It supersedes BAUDIS 99B.
- 115 ARNOLD 99 limit based on the nuclear matrix elements of STAUDT 90.
- 116 BERNATOWICZ 92 finds these majorana neutrino mass limits assuming that the measured geochemical decay width is a limit on the 0ν decay width. The range is the range found using matrix elements from HAXTON 84, TOMODA 87, and SUHONEN 91. Further details of the experiment are given in BERNATOWICZ 93.
- 117 ELLIOTT 92 uses the matrix elements of HAXTON 84.

NODE=S076MW;LINKAGE=RG

NODE=S076MW;LINKAGE=AG

NODE=S076MW;LINKAGE=AL

NODE=S076MW;LINKAGE=UE

NODE=S076MW;LINKAGE=NA

NODE=S076MW;LINKAGE=NR

NODE=S076MW;LINKAGE=KR

NODE=S076MW;LINKAGE=AO

NODE=S076MW;LINKAGE=N1

NODE=S076MW;LINKAGE=N2

NODE=S076MW;LINKAGE=AN

NODE=S076MW;LINKAGE=A1

NODE=S076MW;LINKAGE=KL

NODE=S076MW;LINKAGE=OG

NODE=S076MW;LINKAGE=RR

NODE=S076MW;LINKAGE=VF

NODE=S076MW;LINKAGE=SH

NODE=S076MW;LINKAGE=D5

NODE=S076MW;LINKAGE=KX

NODE=S076MW;LINKAGE=EJ

NODE=S076MW;LINKAGE=K1

NODE=S076MW;LINKAGE=LD

NODE=S076MW;LINKAGE=BB

NODE=S076MW;LINKAGE=L

Limits on Lepton-Number Violating ($V+A$) Current Admixture

For reasons given in the discussion at the beginning of this section, we list only results from 1989 and later. $\langle \lambda \rangle = \lambda \sum U_{ej} V_{ej}$ and $\langle \eta \rangle = \eta \sum U_{ej} V_{ej}$, where the sum is over the number of neutrino generations. This sum vanishes for massless or unmixed neutrinos. In the following Listings, only best or comparable limits or lifetimes for each isotope are reported.

$\langle \lambda \rangle (10^{-6})$	CL%	$\langle \eta \rangle (10^{-8})$	CL%	ISOTOPE	METHOD	DOCUMENT ID
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<120	90			¹⁰⁰ Mo	$0^+ \rightarrow 2^+$	118 ARNOLD 07
$0.692^{+0.058}_{-0.056}$	68	$0.305^{+0.026}_{-0.025}$	68	⁷⁶ Ge	Enriched HPGe	119 KLAUDOR-K... 06A
< 2.5	90			¹⁰⁰ Mo	0ν , NEMO-3	120 ARNOLD 05A
< 3.8	90			⁸² Se	0ν , NEMO-3	121 ARNOLD 05A
< 1.5–2.0	90			¹⁰⁰ Mo	0ν , NEMO-3	122 ARNOLD 04
< 3.2–3.8	90			⁸² Se	0ν , NEMO-3	123 ARNOLD 04
< 1.6–2.4	90	< 0.9–5.3	90	¹³⁰ Te	Cryog. det.	124 ARNABOLDI 03
< 2.2	90	<2.5	90	¹¹⁶ Cd	¹¹⁶ CdWO ₄ scint.	125 DANEVICH 03
< 3.2–4.7	90	< 2.4–2.7	90	¹⁰⁰ Mo	ELEGANT V	126 EJIRI 01
< 1.1	90	<0.64	90	⁷⁶ Ge	Enriched HPGe	127 GUENTHER 97
< 4.4	90	<2.3	90	¹³⁶ Xe	TPC	128 VUILLEUMIER 93
		<5.3		¹²⁸ Te	Geochem	129 BERNATOWICZ 92

118 ARNOLD 07 use NEMO-3 half life limit for 0ν -decay of ¹⁰⁰Mo to the first excited 2^+ -state of daughter nucleus to limit the right-right handed admixture of weak currents $\langle \lambda \rangle$. This limit is not competitive when compared to the decay to the ground state.

119 Re-analysis of data originally published in KLAUDOR-KLEINGROTHAUS 04A. Modified pulse shape analysis leads the authors to claim 6σ statistical evidence for observation of 0ν -decay. Authors use matrix element of MUTO 89 to determine $\langle \lambda \rangle$ and $\langle \eta \rangle$. Uncertainty of nuclear matrix element is not reflected in stated errors.

120 ARNOLD 05A derive limit for $\langle \lambda \rangle$ based on ¹⁰⁰Mo data collected with NEMO-3 detector. No limit for $\langle \eta \rangle$ is given. Supersedes ARNOLD 04.

121 ARNOLD 05A derive limit for $\langle \lambda \rangle$ based on ⁸²Se data collected with NEMO-3 detector. No limit for $\langle \eta \rangle$ is given. Supersedes ARNOLD 04.

122 ARNOLD 04 use the matrix elements of SUHONEN 94 to obtain a limit for $\langle \lambda \rangle$, no limit for $\langle \eta \rangle$ is given. This limit is more stringent than the limit in EJIRI 01 for the same nucleus.

123 ARNOLD 04 use the matrix elements of TOMODA 91 and SUHONEN 91 to obtain a limit for $\langle \lambda \rangle$, no limit for $\langle \eta \rangle$ is given.

124 Supersedes ALESSANDRELLO 00. Cryogenic calorimeter search. Reported a range reflecting uncertainty in nuclear matrix element calculations.

125 Limits for $\langle \lambda \rangle$ and $\langle \eta \rangle$ are based on nuclear matrix elements of STAUDT 90. Supersedes DANEVICH 00.

126 The range of the reported $\langle \lambda \rangle$ and $\langle \eta \rangle$ values reflects the spread of the nuclear matrix elements. On axis value assuming $\langle m_\nu \rangle = 0$ and $\langle \lambda \rangle = \langle \eta \rangle = 0$, respectively.

127 GUENTHER 97 limits use the matrix elements of STAUDT 90. Supersedes BALYSH 95 and BALYSH 92.

128 VUILLEUMIER 93 uses the matrix elements of MUTO 89. Based on a half-life limit 2.6×10^{23} y at 90%CL.

129 BERNATOWICZ 92 takes the measured geochemical decay width as a limit on the 0ν width, and uses the SUHONEN 91 coefficients to obtain the least restrictive limit on η . Further details of the experiment are given in BERNATOWICZ 93.

NODE=S076ETA

NODE=S076ETA

NODE=S076ETA

OCCUR=2

OCCUR=2

NODE=S076ETA;LINKAGE=NA

NODE=S076ETA;LINKAGE=KR

NODE=S076ETA;LINKAGE=N1

NODE=S076ETA;LINKAGE=N2

NODE=S076ETA;LINKAGE=A1

NODE=S076ETA;LINKAGE=A2

NODE=S076ETA;LINKAGE=RR

NODE=S076ETA;LINKAGE=VG

NODE=S076ETA;LINKAGE=EJ

NODE=S076ETA;LINKAGE=J1

NODE=S076ETA;LINKAGE=V

NODE=S076ETA;LINKAGE=BB

NODE=S076

REFID=54331

REFID=54257

REFID=54328

REFID=54483

REFID=54083

REFID=16661

REFID=16385

REFID=16389

REFID=53862

REFID=16373

REFID=53491

REFID=53385

REFID=53053

REFID=52900

REFID=52783

REFID=52519

REFID=52084

REFID=52497

REFID=52522

REFID=51598

REFID=51770

REFID=51401

REFID=50930

Double- β Decay REFERENCES

ANDREOTTI	12	PR C85 045503	E. Andreotti <i>et al.</i>	(CUORICINO Collab.)
AUGER	12	PRL 109 032505	M. Auger <i>et al.</i>	(EXO-200 Collab.)
BELLI	12A	PR C85 044610	P. Belli <i>et al.</i>	
GANDO	12A	PR C85 045504	A. Gando <i>et al.</i>	(KamLAND-Zen Collab.)
ACKERMAN	11	PRL 107 212501	N. Ackerman <i>et al.</i>	(EXO Collab.)
ARNOLD	11	PRL 107 062504	R. Arnold <i>et al.</i>	(NEMO-3 Collab.)
BARABASH	11	PR C83 045503	A.S. Barabash <i>et al.</i>	
BARABASH	11A	PAN 74 312	A.S. Barabash <i>et al.</i>	(NEMO-3 Collab.)
		Translated from YAF 74 330.		
BELLI	11D	JPG 38 115107	P. Belli <i>et al.</i>	(DAMA-INR Collab.)
RUKHADZE	11	NP A852 197	N.I. Rukhadze <i>et al.</i>	(TGV-2 Collab.)
ARGYRIADES	10	NP A847 168	J. Argyriades <i>et al.</i>	(NEMO-3 Collab.)
BELLI	10	NP A846 143	P. Belli <i>et al.</i>	(DAMA-INR Collab.)
ARGYRIADES	09	PR C80 032501	J. Argyriades <i>et al.</i>	(NEMO-3 Collab.)
BELLI	09A	NP A826 256	P. Belli <i>et al.</i>	(DAMA-INR Collab.)
KIDD	09	NP A821 251	M. Kidd <i>et al.</i>	
ARNABOLDI	08	PR C78 035502	C. Arnaboldi <i>et al.</i>	
BELLI	08	PL B658 193	P. Belli <i>et al.</i>	(DAMA-INR Collab.)
BELLI	08B	EPJ A36 167	P. Belli <i>et al.</i>	
UMEHARA	08	PR C78 058501	S. Umehara <i>et al.</i>	
ARNOLD	07	NP A781 209	R. Arnold <i>et al.</i>	(NEMO-3 Collab.)
BARABASH	07	NP A785 371	A.S. Barabash <i>et al.</i>	
KLAUDOR-K...	06A	MPL A21 1547	H.V. Klapdor-Kleingrothaus, I.V. Krivosheina	
ARNABOLDI	05	PRL 95 142501	C. Arnaboldi <i>et al.</i>	(CUORICINO Collab.)

NODE=S076

ARNOLD	05A	PRL 95 182302	R. Arnold <i>et al.</i>	(NEMO-3 Collab.)	REFID=50934
AALSETH	04	PR D70 078302	C.E. Aalseth <i>et al.</i>		REFID=50235
ARNABOLDI	04	PL B584 260	C. Arnaboldi <i>et al.</i>		REFID=49824
ARNOLD	04	JETPL 80 377	R. Arnold <i>et al.</i>	(NEMO3 Detector Collab.)	REFID=50411
		Translated from ZETFP 80 429.			
BARABASH	04	JETPL 79 10	A.S. Barabash <i>et al.</i>		REFID=50551
KLAPDOR-K...	04A	PL B586 198	H.V. Klapdor-Kleingrothaus <i>et al.</i>		REFID=49889
KLAPDOR-K...	04B	PR D70 078301	H.V. Klapdor-Kleingrothaus, A. Dietz, I.V. Krivosheina		REFID=50234;ERROR=1
OGAWA	04	NP A730 215	I. Ogawa <i>et al.</i>		REFID=49796
ARNABOLDI	03	PL B557 167	C. Arnaboldi <i>et al.</i>		REFID=49258
CIVITARESE	03	NP A729 867	O. Civitarese, J. Suhonen		REFID=49794
DANEVICH	03	PR C68 035501	F.A. Danovich <i>et al.</i>		REFID=49600
DOERR	03	NIM A513 596	C. Doerr, H.V. Klapdor-Kleingrothaus		REFID=50553
AALSETH	02B	PR D65 092007	C.E. Aalseth <i>et al.</i>	(IGEX Collab.)	REFID=48793
BERNABEI	02D	PL B546 23	R. Bernabei <i>et al.</i>	(DAMA Collab.)	REFID=48990
KLAPDOR-K...	02B	PPNL 110 57	H.V. Klapdor-Kleingrothaus, A. Dietz, I.V. Krivosheina		REFID=49114;ERROR=2
KLAPDOR-K...	02D	FP 32 1181	H.V. Klapdor-Kleingrothaus, A. Dietz, I.V. Krivosheina		REFID=49193;ERROR=3
SIMKOVIC	02	hep-ph/0204278	F. Simkovic, P. Domin, A. Faessler		REFID=49198
ASHITKOV	01	JETPL 74 529	V.D. Ashitkov <i>et al.</i>		REFID=48730
		Translated from ZETFP 74 601.			
DANEVICH	01	NP A694 375	F.A. Danovich <i>et al.</i>		REFID=48113
DEBRAECKEL...	01	PRL 86 3510	L. De Braeckeleer <i>et al.</i>		REFID=48467
EJIRI	01	PR C63 065501	H. Ejiri <i>et al.</i>		REFID=48163
KLAPDOR-K...	01	EPJ A12 147	H.V. Klapdor-Kleingrothaus <i>et al.</i>		REFID=48460
KLAPDOR-K...	01B	MPL A16 2409	H.V. Klapdor-Kleingrothaus <i>et al.</i>		REFID=48628
STOICA	01	NP A694 269	S. Stoica, H.V. Klapdor-Kleingrothaus		REFID=50526
WIESER	01	PR C64 024308	M.E. Wieser, J.R. De Laeter		REFID=48224
ALESSAND...	00	PL B486 13	A. Alessandrello <i>et al.</i>		REFID=47726
BRUDANIN	00	PL B495 63	V.B. Brudanin <i>et al.</i>	(TGV Collab.)	REFID=47917
DANEVICH	00	PR C62 045501	F.A. Danovich <i>et al.</i>		REFID=47810
ARNOLD	99	NP A658 299	R. Arnold <i>et al.</i>	(NEMO Collab.)	REFID=47325
BAUDIS	99	PR D59 022001	L. Baudis <i>et al.</i>	(Heidelberg-Moscow Collab.)	REFID=46702
BAUDIS	99B	PRL 83 41	L. Baudis <i>et al.</i>	(Heidelberg-Moscow Collab.)	REFID=47031
SIMKOVIC	99	PR C60 055502	F. Simkovic <i>et al.</i>		REFID=50525
ARNOLD	98	NP A636 209	R. Arnold <i>et al.</i>	(NEMO-2 Collab.)	REFID=49250
ALSTON-...	97	PR C55 474	M. Alston-Garnjost <i>et al.</i>	(LBL, MTHO+)	REFID=45230
DESILVA	97	PR C56 2451	A. de Silva <i>et al.</i>	(UCI)	REFID=45865
GUENTHER	97	PR D55 54	M. Gunther <i>et al.</i>	(Heidelberg-Moscow Collab.)	REFID=45265
ARNOLD	96	ZPHY C72 239	R. Arnold <i>et al.</i>	(BCEN, CAEN, JINR+)	REFID=45277
BALYSH	96	PRL 77 5186	A. Balysh <i>et al.</i>	(KIAE, UCI, CIT)	REFID=45405
TAKAOKA	96	PR C53 1557	N. Takaoka, Y. Motomura, K. Nagao	(KYUSH, OKAY)	REFID=45407
ARNOLD	95	JETPL 61 170	R.G. Arnold <i>et al.</i>	(NEMO Collab.)	REFID=44332
		Translated from ZETFP 61 168.			
BALYSH	95	PL B356 450	A. Balysh <i>et al.</i>	(Heidelberg-Moscow Collab.)	REFID=44401
BARABASH	95	PL B345 408	A.S. Barabash <i>et al.</i>	(ITEP, SCUC, PNL+)	REFID=44334
DASSIE	95	PR D51 2090	D. Dassie <i>et al.</i>	(NEMO Collab.)	REFID=44148
EJIRI	95	JPSJ 64 339	H. Ejiri <i>et al.</i>	(OSAK, KIEV)	REFID=44333
KOBAYASHI	95	NP A586 457	M. Kobayashi, M. Kobayashi	(KEK, SAGA)	REFID=44175
SUHONEN	94	PR C49 3055	J. Suhonen, O. Civitarese		REFID=50527
ARTEMEV	93	JETPL 58 262	V.A. Artemiev <i>et al.</i>	(ITEP, INRM)	REFID=44331
		Translated from ZETFP 58 256.			
BERNATOW...	93	PR C47 806	T. Bernatowicz <i>et al.</i>	(WUSL, TATA)	REFID=43219
KAWASHIMA	93	PR C47 R2452	A. Kawashima, K. Takahashi, A. Masuda	(TOKYC+)	REFID=44329
VUILLEUMIER	93	PR D48 1009	J.C. Vuilleumier <i>et al.</i>	(NEUC, CIT, VILL)	REFID=43446
BALYSH	92	PL B283 32	A. Balysh <i>et al.</i>	(MPIH, KIAE, SASSO)	REFID=42083
BERNATOW...	92	PRL 69 2341	T. Bernatowicz <i>et al.</i>	(WUSL, TATA)	REFID=42181
BLUM	92	PL B275 506	D. Blum <i>et al.</i>	(NEMO Collab.)	REFID=41979
ELLIOTT	92	PR C46 1535	S.R. Elliott <i>et al.</i>	(UCI)	REFID=42190
EJIRI	91	PL B258 17	H. Ejiri <i>et al.</i>	(OSAK)	REFID=41483
MANUEL	91	JPG 17 S221	O.K. Manuel	(MISSR)	REFID=45408
SUHONEN	91	NP A535 509	J. Suhonen, S.B. Khadkikar, A. Faessler	(JYV+)	REFID=41945
TOMODA	91	RPP 54 53	T. Tomoda		REFID=45867
TURKEVICH	91	PRL 67 3211	A. Turkevich, T.E. Economou, G.A. Cowan	(CHIC+)	REFID=41832
YOU	91	PL B265 53	K. You <i>et al.</i>	(BHEP, CAST+)	REFID=41598
STAUDT	90	EPL 13 31	A. Staudt, K. Muto, H.V. Klapdor-Kleingrothaus		REFID=41834;ERROR=4
MUTO	89	ZPHY A334 187	K. Muto, E. Bender, H.V. Klapdor	(TINT, MPIH)	REFID=43301
LIN	88	NP A481 477	W.J. Lin <i>et al.</i>		REFID=49526
LIN	88B	NP A481 484	W.J. Lin <i>et al.</i>		REFID=49527
BOEHM	87	Massive Neutrinos Cambridge Univ.	F. Boehm, P. Vogel	(CIT)	REFID=41833
TOMODA	87	PL B199 475	T. Tomoda, A. Faessler	(TUBIN)	REFID=43303
HAXTON	84	PPNP 12 409	W.C. Haxton, G.J. Stevenson		REFID=43302
KIRSTEN	83	PRL 50 474	T. Kirsten, H. Richter, E. Jessberger	(MPIH)	REFID=10425